

Duality in strong interactions^{*}

L. Jenkovszky^{1*}, V.K. Magas^{1,2†} and E. Predazzi^{3◇}

¹ *Bogolyubov Institute for Theoretical Physics
Academy of Sciences of Ukraine
01143 Kiev, Ukraine*

² *Center for Physics of Fundamental Interactions
Instituto Superior Tecnico
Av. Rovisco Pais, 1049-001 Lisbon, Portugal*

³ *Torino University and INFN
via P. Giuria, 1, Torino, 10125 Italy*

^{*} *Email: jenk@gluk.org*

[†] *Email: vladimir@cfif.ist.utl.pt*

[◇] *Email: predazzi@to.infn.it*

February 8, 2008

In this lecture we first present the classical results on duality in the strong interactions, developed in the late 60-es and the early 70-es, regarding on-mass-shell hadronic reactions in the framework of the analytical S -matrix theory. Then we discuss their extension to off-mass-shell processes (virtual Compton- and deep inelastic scattering), published in a series of recent papers [1, 2, 3]

The continuity of dynamics in going from low to high energies, from low to high virtualities, from exclusive to inclusive processes etc is a basic principle of the strong interactions (see [4]). Duality between Reggeon-resonance is a particularly nice and instructive example showing how it

^{*}Lecture delivered by L. Jenkovszky at the 6th International Summer School-Seminar On Actual Problems Of High Energy Physics, Gomel, Belarus, August 7-16, 2001.

works. Before its discovery, the contribution from low-energy, direct channel resonances was considered to be independent from that of high-energy Regge exchanges, and the complete scattering amplitude was thought to be their sum. This was the interference model. Later on, in analyzing finite energy sum rules, it was realized that each piece already contains the whole dynamics and that a proper sum of direct channel resonances "knows" its high energy, Regge behavior and vice versa: from a smooth Regge asymptotic formula resonances can be created and thus the sum of s -channel resonances and t -channel exchanges (interference model) leads to wrong "double counting". Instead, their equivalence, or duality proved to be the right answer. In a similar way, below we shall ask in which way the small- and large- x parts of the structure functions are related.

An important further step in the late 60-es consisted in the construction of explicit models realizing duality. The Veneziano model [5] is the best known example. The common feature of most of the dual models was their "soft" dynamics, which means that they were applicable only for peripheral ("soft") collisions and did not contain any information on the interaction at short distances, where the partonic nature of the strong interaction may reveal. The reason for this shortcoming was in the linear character of the Regge trajectories admitted in those "narrow resonance" dual models. Their contradiction with analyticity and unitarity and the absence of any imaginary part was the main reason why the physical applications of dual models were abandoned, giving place to formal mathematical developments.

A revival of this subject was partly due to the recent experimental measurements of the nucleon structure function at JLab (CEBAF) [6]. They inspired the present authors in the construction of a unified "two-dimensionally dual" picture of the strong interaction [1, 2, 3] connecting low- and high energies (Veneziano, or resonance-reggeon duality [5]) with low- and high virtualities (Q^2) (Bloom-Gilman, or hadron-parton duality [7]). The basic idea of the unification is the use of a Q^2 -dependent dual amplitudes, employing nonlinear complex Regge trajectories providing an imaginary part in the scattering amplitude, related to the total cross section and structure functions and thus saturating duality by a finite number of resonances lying on the (limited) real part of the Regge trajectories.

The resulting object, a deeply virtual scattering amplitude, $A(s, t, Q^2)$, is a function of three variables, reducing to a nuclear structure function (SF) when $t = 0$ and to an on-shell hadronic scattering amplitude for

$Q^2 = m^2$. It closes the circle in Fig 1. We use this amplitude to describe the background as well as the resonance component [8].




Figure 1: Veneziano, or resonance-reggeon duality [5] and Bloom-Gilman, or hadron-parton duality [7] in strong interactions

The Q^2 -dependence of the residue functions here will be chosen in such a way as to provide for Bjorken scaling at small x (large s). The resulting amplitude (structure function) is applicable in the whole kinematical range, including the resonance region. We call this unification "two dimensional duality" - one in s , the other one in Q^2 ,

In the early days of duality, off mass continuation was attempted [9] by means of multi-leg (e.g. 6-point) dual amplitudes with "extra" lines taken at their poles. Without going into details, here we only mention that scaling in this approach can be achieved [10] only with nonlinear trajectories, e.g. those with logarithmic or constant asymptotic.

1 Notation and conventions

We use standard notation for the cross section and structure function (see Fig. 2):

$$\sigma^{\gamma^*p} = \frac{4\pi^2\alpha(1 + 4m^2x^2/Q^2)}{Q^2(1 - x)} \frac{F_2(x, Q^2)}{1 + R(x, Q^2)}, \quad (1)$$

where α is the fine structure constant, Q^2 is minus the squared four momentum transfer or the momentum carried by the virtual photon, x is the Bjorken variable and s is the squared center of mass energy of the γ^*p system, obeying the relation

$$s = Q^2(1 - x)/x + m_p^2, \quad (2)$$

where m_p is the proton mass and $R(x, Q^2) = \sigma_L(x, Q^2)/\sigma_T(x, Q^2)$. For the sake of simplicity we set $R = 0$, which is a reasonable approximation.




Figure 2: Kinematic of deep inelastic scattering.

We use the norm where

$$\sigma_T^{\gamma^*}(s, t, Q^2) = \text{Im } A(s, t, Q^2). \quad (3)$$

According to the two-component duality picture [8] both the scattering amplitude A and the structure function F_2 are sums of a diffractive and non-diffractive terms. At high energies both terms are Regge-behaved. In γ^*p scattering only positive signature exchange are allowed. The dominant ones are the Pomeron and the f Reggeon, respectively. The relevant scattering amplitude is (remember that here $t = 0$)

$$A_i(s, Q^2) = i\beta_k(Q^2) \left(-i \frac{s}{s_i} \right)^{\alpha_k(0)-1}, \quad (4)$$

where α and β are the Regge trajectory and residue and k stands either for the Pomeron or the Reggeon. As usual, the residue will be chosen such as to satisfy approximate Bjorken scaling for the structure function [22, 23]. It should be remembered that by factorization, assuming that the Reggeon (or Pomeron) exchange is a simple pole, the residue function is a product of two vertices - the $\gamma\gamma R(P)$ and $NNR(P)$, where N stands for the nucleon (see Fig. 3).

At low energies the scattering amplitude is dominated by the contribution of the near-by resonances. In the vicinity of a resonance Res , the amplitude can be also written in a factorized form, product of the probabilities that two particles - γ and p form a resonance with squared mass s_R and total width Γ

$$A(s, Q^2) = \sum_{spin} \frac{A_{fi}(Q^2)A_{if}^*(Q^2)}{s_{Res} - s - i\Gamma}, \quad (5)$$

where the s_R is the squared mass of the resonance and $\Gamma(Q^2)$ is its width; A_{fi} are the inelastic form factors.




Figure 3: According to the Veneziano (or resonance-reggeon) duality a proper sum of either t-channel or s-channel resonance exchanges accounts for the whole amplitude. *Diagram drawings in Figs. 2 and 3 have been made with **scribble1.2** program [25].*

2 Nucleon resonances in inelastic electron-nucleon scattering

Some thirty years ago Bloom and Gilman [7] observed that the prominent resonances in inelastic electron-proton scattering do not disappear with increasing Q^2 relatively to the "background" but instead fall at roughly the same rate as any background. Furthermore, the smooth scaling limit

proved to be an accurate average over resonance bumps seen at lower Q^2 and s .

Since then, the phenomenon was studied in a number of papers [11, 12, 13] and recently has been confirmed experimentally [6] (see Fig. 4). These studies were aimed mainly to answer the questions: in which way a limited number of resonances can reproduce the smooth scaling behavior? The main theoretical tools in these studies were finite energy sum rule and perturbative QCD calculations, whenever applicable. Our aim instead is the construction of an explicit dual model combining direct channel resonances, Regge behavior, typical of hadrons and scaling behavior, typical of the partonic picture.

The existence of resonances in the structure function at large x , close to $x = 1$ by itself is not surprising: by the relations (1) and (2) they are the same as in γ^*p total cross section, seen in different variables. The important question is if and how a small number of resonances (or even a single one) can reproduce the smooth Bjorken scaling behavior, known to be an asymptotic property, typical of multiparticle processes.

The possibility that a limited (small) number of resonances can build up smooth Regge behavior was demonstrated by means of finite energy sum rules [14]. Further it was confused by the presence of an infinite number narrow resonances in the Veneziano model [5], which made its phenomenological application difficult, if not impossible. Similar to the case of the resonance-reggeon duality [14], hadron-parton duality was established [7] by means of the finite energy sum rules, but it was not realized explicitly like the Veneziano model (or its further modifications).

Actually, the early onset of Bjorken scaling, called "early, or precaution scaling" was observed with the first measurements of deep inelastic scattering at SLAC, where it was noticed that a more rapid approach to scaling can be achieved with the BG variable [7] $x' = x/(1 + m^2x^2/Q^2)$ instead of x (or $\omega = 1/x$). More recently the following generalization of the BG variable

$$\xi = \frac{2x}{1 + \sqrt{1 + \frac{4m^2x^2}{Q^2}}} \quad (6)$$

was suggested by O.Nachtmann [16]. We use the standard Bjorken variable x , however our results can be easily rewritten in terms of the above-mentioned modified variables.

First attempts to combine resonance (Regge) behavior with Bjorken scaling were made [15, 17, 18] at low energies (large x), with the emphasis




Figure 4: Extracted νW_2 structure function spectra as a function of Nachtmann scaling variable (6). The solid curve indicates a fit to DIS data for $Q^2 = 10 \text{ GeV}^2$, dashed curve - $Q^2 = 5 \text{ GeV}^2$. From [6].

on the right choice of the Q^2 -dependence, such as to satisfy the required behavior of the form factors, vector meson dominance (VMD) with the requirement of the Bjorken scaling. (N.B.: the validity (or failure) of the (generalized) VMD is still disputed). Similar attempts in the high-energy (low x) region became popular recently, with the advent of the HERA data. They will be presented in Sec. 5.

A consistent treatment of the problem requires the account for the spin dependence. For simplicity we ignore it in this lecture (see e.g. [13]).

3 Factorization and dual properties (bootstrap) of the vertices

Let us remind that the residue functions are completely arbitrary in the Regge pole model, but they are constrained in the dual model. We show this by using the low energy- (resonances) and high energy- (Regge) decomposition on the simple Veneziano model [5]

$$V(s, t) = \int_0^1 dz z^{-\alpha(s)} (1-z)^{-\alpha(t)} = B(1-\alpha(s), 1-\alpha(t)) = \frac{\Gamma(1-\alpha(s))\Gamma(1-\alpha(t))}{\Gamma(2-\alpha(s)-\alpha(t))}. \quad (7)$$

$$V(s, t) = \sum_{n=1}^{\infty} \frac{1}{n-\alpha(s)} \frac{\Gamma(n+\alpha(t)+1)}{n! \Gamma(\alpha(t)+1)}. \quad (8)$$

By the Stirling formula

$$V(s, t) \Big|_{|\alpha(s)| \rightarrow \infty} \rightarrow [-\alpha(s)]^{\alpha(t)-1} \Gamma(1-\alpha(t)) \left[\sum_{n=0}^N \frac{a_n(0)}{[\alpha(s)]^n} + 0 \left(\frac{1}{[\alpha(s)]^{N+1}} \right) \right] \quad (9)$$

and since for small $|t|$ the Γ function varies slowly compared with the exponential, the Regge asymptotic behavior is

$$V(s, t) \sim (\alpha' s)^{\alpha(t)}, \quad (10)$$

where $\beta(t) = (\alpha')^{\alpha(t)}$ is the Regge residue.

Actually, one has to identify a single (and hence economic!) Regge exchange amplitude with a sum of direct channel poles. Such an identification is not practical for an infinite number of poles (e.g. the Veneziano amplitude) but, as we show below is feasible if their number is finite (small). To anticipate the forthcoming discussion, we shall feed the Q^2 -dependence in the Regge residue at high energies (small x and use the dual amplitude with a finite number of resonances!) to the whole kinematical region, including that of resonances. Relating the amplitude to the SF, we set $t = 0$.

To remedy the problems of the infinite number of narrow resonance, nonunitarity and lack of an imaginary part, we use a generalization of the Veneziano model free from the above-mentioned difficulties.

4 Dual amplitude with Mandelstam analyticity

The invariant dual on-shell scattering amplitude dual amplitude with Mandelstam analyticity (DAMA) applicable both to the diffractive and non-diffractive components reads [19]

$$D(s, t) = \int_0^1 dz \left(\frac{z}{g}\right)^{-\alpha(s')-1} \left(\frac{1-z}{g}\right)^{-\alpha(t')-1}, \quad (11)$$

where $s' = s(1-z)$, $t' = tz$, g is a parameter, $g > 1$, and s , t are the Mandelstam variables.

For $s \rightarrow \infty$ and fixed t it has the following Regge asymptotic behavior

$$D(s, t) \approx \sqrt{\frac{2\pi}{\alpha_t(0)}} g^{1+a+ib} \left(\frac{s\alpha'(0)g \ln g}{\alpha_t(0)}\right)^{\alpha_t(0)-1}, \quad (12)$$

where $a = \text{Re } \alpha\left(\frac{\alpha_t(0)}{\alpha'(0) \ln g}\right)$ and $b = \text{Im } \alpha\left(\frac{\alpha_t(0)}{\alpha'(0) \ln g}\right)$.

The pole structure of DAMA is similar to that of the Veneziano model except that multiple poles may appear at daughter levels. The presence of these multipoles does not contradict the theoretical postulates. On the other hand, they can be removed without any harm to the dual model by means the so-called Van der Corput neutralizer. The procedure [19] is to multiply the integrand of (11) by a function $\phi(x)$ with the properties:

$$\phi(0) = 0, \quad \phi(1) = 1, \quad \phi^n(1) = 0, \quad n = 1, 2, 3, \dots$$

The function $\phi(x) = 1 - \exp\left(-\frac{x}{1-x}\right)$, for example, satisfies the above conditions and results [19] in a standard, "Veneziano-like" pole structure:

$$D(s, t) = \sum_n g^{n+\alpha_t(0)} \frac{C_n}{n - \alpha(s)}, \quad (13)$$

where

$$C_n = \frac{\alpha_t(0)(\alpha_t(0) + 1) \dots (\alpha_t(0) + n + 1)}{n!}. \quad (14)$$

The pole term (13) is a generalization of the Breit-Wigner formula (5), comprising a whole sequence of resonances lying on a complex trajectory

$\alpha(s)$. Such a "reggeized" Breit-Wigner formula has little practical use in the case of linear trajectories, resulting in an infinite sequence of poles, but it becomes a powerful tool if complex trajectories with a limited real part and hence a restricted number of resonances are used. Moreover, it appears that a small number of resonances is sufficient to saturate the direct channel.

Contrary to the Veneziano model, DAMA (11) not only allows but rather requires the use of nonlinear complex trajectories providing, in particular, for the imaginary part of the amplitude, resonance widths and resulting in a finite number of those. More specifically, the asymptotic rise of the trajectories in DAMA is limited by the condition (in accordance with an important upper bound derived earlier [20])

$$\left| \frac{\alpha(s)}{\sqrt{s} \ln s} \right| \leq \text{const}, \quad s \rightarrow \infty. \quad (15)$$

Actually, this upper bound can be even lowered up to a logarithm by requiring wide angle scaling behaviour for the amplitude.

Models of Regge trajectories combining the correct threshold and asymptotic behaviors have been widely discussed in the literature (see e.g. [21] for a recent treatment of this problem). A particularly simple model is based on a sum of square roots

$$\alpha(s) = \alpha_0 + \sum_i \gamma_i \sqrt{s_i - s},$$

where the lightest threshold (made of two pions or a pion and a nucleon) is important for the imaginary part, while the heaviest threshold limits the rise of the real part, where resonances terminate.

DAMA with the trajectories specified above is equally applicable to both: the diffractive and non-diffractive components of the amplitude, the difference being qualitative rather than quantitative. The utilization of a trajectory with a single threshold,

$$\alpha_E(s) = \alpha_E(0) + \alpha_{1E}(\sqrt{s_E} - \sqrt{s_E - s}) \quad (16)$$

prevents the production of resonances on the the physical sheet, although they are present on the nonphysical sheet, sustaining duality (i.e. their sum produces Regge asymptotic behavior). This nontrivial property of DAMA makes it particularly attractive in describing the smooth background (dual

to the Pomeron exchange) (see [19]). The threshold value, slope and the intercept of this exotic trajectory are free parameters.

For the resonance component a finite sum in (13) is adequate, but we shall use a simple model with lowest threshold included explicitly and the higher ones approximated by a linear term:

$$\alpha_R(s) = \alpha_R(0) + \alpha' s + \alpha_{1R}(\sqrt{s_0} - \sqrt{s_0 - s}), \quad (17)$$

where s_0 is the lowest threshold – $s_0 = (m_\pi + m_p)^2$ in our case – while the remaining 3 parameters will be adjusted to the known properties of the relevant trajectories (those of N^* and the Δ isobar in our case). The termination of resonances, provided in DAMA by the limited real part, here will be effectively taken into account by a cutoff in the summation of (13).

Finally, we note that a minimal model for the scattering amplitude is a sum

$$A(s, t, u) = c(D(s, t) + D(u, t)), \quad (18)$$

providing the correct signature at high-energy limit, c is a normalization factor. We disregard the symmetry (spin and isospin) properties of the problem, concentrating on its dynamics. In the limit $s \rightarrow \infty$, $t = 0$ we have $u = -s$ and therefore

$$A(s, 0, -s)|_{s \rightarrow \infty} = c D(s, 0)(1 + (-1)^{\alpha_t(0)-1}), \quad (19)$$

where $D(s, t)$ is given by eq. (12). For the total cross section in this limit we obtain:

$$\sigma_T^* = \text{Im } A = C g^{\alpha_t(0)+a} (s \alpha'(0) \ln g)^{\alpha_t(0)-1} \cdot (\sin(\alpha_t(0) - 1)\pi \cos(b \ln g) + (1 + \cos(\alpha_t(0) - 1)\pi) \sin(b \ln g)) , \quad (20)$$

where C is a constant independent on s , g and $\alpha'(0)$.

5 Q^2 – dependence

Our main idea is the introduction of the Q^2 -dependence in the dual model by matching its Regge asymptotic behavior and pole structure to standard forms, known from the literature. The point is that the correct identification of this Q^2 -dependence in a single asymptotic limit of the dual amplitude will extend it to the rest of the kinematical regions. We have

two ways to do so:

A) Combine Regge behavior and Bjorken scaling limits of the structure functions (or Q^2 -dependent γ^*p cross sections)

B) Introduce properly Q^2 dependence in the resonance region.

They should match if the procedure is correct and the dual amplitude should take care of any further inter- or extrapolation.




Figure 5: $g(Q^2)$ - the solution of the transcendent equation (22) - for Δ and exotic trajectories.

It is obvious from eq. (4) that asymptotic Regge and scaling behavior require the residue to fall like $\sim (Q^2)^{-\alpha_i(0)+1}$. Actually, it could be more involved if we require the correct $Q^2 \rightarrow 0$ limit to be respected and the observed scaling violation (the "HERA effect") to be included. Various models to cope with the above requirements have been suggested [22, 23, 24]. At HERA, especially at large Q^2 , scaling is so badly violated that it may not be explicit anymore.

In combining Regge asymptotic behavior with (approximate) Bjorken scaling, one can proceed basically in the following ways – keep explicitly a scaling factor x^Δ (to be broken by some Q^2 -dependence "properly" taken into account) [23]

$$F_2(x, Q^2) \sim x^{-\Delta(Q^2)} \left(\frac{Q^2}{Q^2 + Q_0^2} \right)^{1+\Delta(Q^2)}, \quad (21)$$

where $\Delta(Q^2) = \alpha_t(0) - 1$ may be a constant, in particular.

In the Regge asymptotic limit of the Veneziano model - $\sim (-\alpha' s)^{\alpha(t)-1}$, - Q^2 -dependence can be introduced only through the slope of the direct channel trajectory [1], i.e. by making α'_s Q^2 - dependent. This is a crucial point to be understood. By Regge-factorization, the trajectory should not depend on the parameters (masses, virtualities) of the external particles. On the other hand, there is a nice bootstrap condition in dual models, by which one of the states on the trajectory (e.g. the lightest one) should be identified with the external particle, which means that the slope is associated with its squared mass. This can be generalized to a varying mass or Q^2 .

A different interpretation may come from treating a Regge pole as and effective one, accounting for unitarization etc and consequently embodying a complicated Q^2 -dependence. The latter interpretation became popular [22, 23] in the applications of Regge formulae to explain the drastic increase of the structure functions with Q^2 , observed at HERA (Bjorken scaling violation). Although we do not exclude this possibility (treating it as “effective” Regge pole, we study here the different option of introducing scaling violation in the constant g appearing, besides α' , in the the residue of DAMA, eq (11).

From the explicit Regge asymptotic form of DAMA, (20), and neglecting the logarithmic dependence of g we make the following identification

$$g(Q^2)^{\alpha_t(0)+a} = \left(\frac{Q_{lim}^2}{Q^2 + Q_0^2} \right)^{\alpha_t(0)}. \quad (22)$$

Note that eq. (22) is transcendent with respect to g , since $a = a(g) = Re \alpha \left(\frac{\alpha_t(0)}{\alpha'(0) \ln g} \right)$. Another point to mention is that this equation is not valid in the whole range of Q^2 , since for Q^2 close to Q_{lim}^2 , g may get smaller than 1, which is unacceptable in DAMA. For large Q^2 the Q^2 -dependence of the $\log g$ and $b = b(Q^2) = Im \alpha \left(\frac{\alpha_t(0)}{\alpha'(0) \ln g} \right)$ in eq. (20) can not be neglected, it might contribute to scaling violation.

6 Scaling at large x

Let us now consider the extreme case of a single resonance contribution.




Figure 6: γ^*p total cross section as a function of $\sqrt{s}$. The dashed line shows the contribution from the Δ resonance, the dot-dashed line corresponds to the background, i.e. the contribution from the exotic trajectory. Here $Q^2 = 1 \text{ GeV}^2$.

A resonance pole in DAMA contributes with

$$A(s, t) = g^{n+\alpha_t(0)} \frac{C_n}{n - \alpha(s)}.$$

At the resonance $s = s_{Res}$ one has $Re \alpha(s_{Res}) = n$ and $\frac{Q^2(1-x)}{x} = s_{Res} - m^2$, hence

$$F_2(x, Q^2) = \frac{Q^2(1-x)}{4\pi^2\alpha\left(1 + \frac{4m^2x^2}{Q^2}\right)} \frac{C_n}{Im \alpha(s_{Res})} g(Q^2)^{n+\alpha_t(0)}.$$

As $x \rightarrow 1$ $Q^2 \approx \frac{s_{Res}-m^2}{1-x} \rightarrow \infty$ and

$$F_2(x, Q^2) \sim g\left(\frac{s_{Res} - m^2}{1-x}\right)^{n+\alpha_t(0)}.$$

By using the approximate solution $g(Q^2) \approx (Q_{lim}^2/Q^2)^{\frac{\alpha_t(0)}{\alpha_t(0)+a}}$, where a is a slowly varying function of g , we get for x near 1

$$F_2(x, Q^2) \sim (1-x)^{\frac{\alpha_t(0)(n+\alpha_t(0))}{\alpha_t(0)+a}},$$

where the limits for x are defined by $Q_0^2 \ll \frac{s_{Res}-m^2}{1-x} \leq Q_{lim}^2$.

We recognize a typical large- x scaling behavior $(1-x)^N$ with the power N (counting the quarks in the reaction) depending basically on the intercept of the t -channel trajectories.




Figure 7: The same as Fig. 6, but $Q^2 = 6 \text{ GeV}^2$.

7 Numerical Estimates

Having fixed the Q^2 -dependence of the dual model by matching its Regge asymptotic behavior with that of the structure functions, we now use this dual model to extrapolate down to the resonance region, where its pole expansion (13) is appropriate - now complemented with a Q^2 -dependence through $g(Q^2)$, fixed by eq. (22)

Table 1: Values of parameters used in the calculations shown in Figs. 5, 6 and 7.

	Δ Resonance	Background
Q_{lim}^2, GeV^2	62	120
Q_0^2, GeV^2	0.01	2.5
Dual trajectory	$\alpha_f(t)$ is dual to $\alpha_\Delta(s)$	$\alpha_P(t)$ is dual to $\alpha_E(s)$
	$\alpha_f(0) = 0.9$	$\alpha_P(0) = 1 + 0.077 \cdot \left(1 + \frac{2Q^2}{Q^2 + 1.117}\right)$ [23]

Normalization coefficient $c = 0.03$.

As already said, we write the imaginary part of the scattering amplitude as the sum of two terms - a diffractive (background) and non-diffractive (resonance) one. Note that $g(Q^2)$ has the same functional form (22) in both cases, only the values of the parameters differ (they are fixed from the small- x fits [24] of the SF).

At low, resonance, energies γ^*p scattering exhibits a rich resonance structure, intensively studied in a number of papers. About 20 resonances overlap, their relative importance varying with Q^2 , but only a few can be identified more or less unambiguously. These are: $\Delta^+(1236)$ with $J^P = \frac{3}{2}^+$, $N^{*+}(1520)$, $J^P = \frac{3}{2}^-$, $N^{*+}(1688)$, $J^P = \frac{5}{2}^+$ and $N^{*+}(1920)$ with $J^P = \frac{7}{2}^+$ (see Fig. 9). They lie on the Δ and the exchange-degenerate N trajectories. In this work we are mainly interested in introducing Q^2 -dependence into the scattering amplitude, therefore we will concentrate on a single resonance ($\Delta^+(1236)$) at different values of Q^2 . We use trajectories (17) in which the lowest pion-nucleon threshold is included explicitly, while higher thresholds are approximated by a linear term:

$$\alpha_\Delta(s) = 0.1 + 0.84s + 0.1331(\sqrt{s_0} - \sqrt{s_0 - s}),$$

where $s_0 = (m_\pi^2 + m_N^2)$.¹ The above values of the parameters are chosen so as to fit the known mass and width of the Δ resonance in a way consistent with the known linear parameterizations.

¹ Actually, trajectories without any linear term (see e.g. [21]) could be more appropriate (and will be studied in future).




Figure 8: γ^*p total cross section as a function of $\sqrt{s}$ and Q^2 . For different values of Q^2 we show the contributions from the Δ resonance (dashed line), the background, i.e. the contribution from the exotic trajectory (dot-dashed line) and their sum (full line).

In the interval of interest $\sqrt{s} = 1.1 - 1.5 \text{ GeV}$, $t = 0$ we have $u = m_N^2 - s < 0$, so, it is far from resonance region, therefore we neglect the contribution from $D(u, t)$ for both resonance and background terms.

The smooth background is also modeled by a single term and exotic trajectory (16). As already explained, the direct channel Regge pole does not produce here physical resonances. The parameters of the exotic trajectory are the following:

$$\alpha_E(s) = -0.25 + 0.25(\sqrt{1.21} - \sqrt{1.21 - s}), \quad (23)$$

where $s_E = 1.1^2 \text{ GeV}^2$ is an effective exotic threshold. Obviously "pole" doesn't mean a resonance in this case.

Figure 5 shows g as a function of Q^2 for Δ and exotic trajectories. The resulting cross sections (imaginary part of the amplitude) in the resonance

region is shown in Figs. 6 and 7 for two values of $Q^2 = 1$ and 6 GeV^2 . It is in qualitative agreement with the experimental data (compare with Fig. 9) [2, 3]. Figure 8 shows the dual properties of the cross section in 2 dimensions - one is the squared energy s and the other one is virtuality Q^2 . Table 1 shows the values of the parameters used in our calculations.

8 Conclusions

The spirit of the present lectures is close to that of the remarkable paper [4]. Moreover, we are trying to construct an explicit model realizing the continuity of the strong interactions. We have shown how Q^2 dependence can be incorporated in DAMA, becoming a function of 3 variables, namely s, t and Q^2 and thus making complete the circle shown in Fig. 1. In particular, we answer the question posed at the beginning of the present lectures: "Does the small- x structure function know its large- x behavior (and v.v.)?". Our answer is affirmative.

The main conclusions from our analysis are that:

- A) Q^2 -dependence at low- and high- x (or high- and low- s) are interrelated and have the same origin;
- B) Even a single (low energy) resonance can produce a smooth scaling-like curve in the structure function (parton-hadron duality).

To summarize, we have suggested an explicit dual model in which the Q^2 -dependence introduced in the low- x domain is extended to the whole kinematic region, in particular to the region of resonances. The resulting predictions for the first resonance in the γ^*p system shown in Figs. 6, 7 are in quantitative agreement with data.

9 Acknowledgments

L.J. and V.M. acknowledge the support by INTAS, Grant 00-00366. The work of L.J. was supported also by the US Civilian Research and Development Foundation (CRDF), Grant UP1-2119.

L.J. thanks the Organizers of the Gomel School-Seminar for the invitation and for the warm, pleasant and creative atmosphere during the Seminar and Seminar.

References

- [1] L.L. Jenkovszky, V.K. Magas and F. Paccanoni, Proceedings of the "New Trend in High-Energy Physics", Crimea, Ukraine, May 27 - June 4, 2000, p. 121; R. Fiore, L. Jenkovszky, V. Magas, Proceedings of the "Diffractin-2000", Cetraro, Italy, 2-8 September 2000, Nucl. Phys. B (Proceedings Supplements) **99**, (2001) 131.
- [2] L. Jenkovszky and V. Magas, to be published in the Proceedings of the 9th Blois Workshop On Elastic and Diffractive Scattering, June 9-15, 2001, Prague, Czech Republic; to be published in the Proceedings of "Spin-2001", 9th International Workshop On High-Energy Spin Physics, 2-7 Aug 2001, Dubna, Russia; to be published in the Proceedings of the "ISMD 2001", XXXI International Symposium on Multiparticle Dynamics, September 1-7, 2001, Datong, China; L. Jenkovszky, T. Korzhinskaya and V. Magas, to be published in the Proceedings of the "New Trend in High-Energy Physics", Yalta, Crimea, Ukraine, Sept 22-29, 2001.
- [3] L.L. Jenkovszky, V.K. Magas and E. Predazzi, hep-ph/0110374.
- [4] J.D. Bjorken, J.B. Kogut Phys. Rev. **D 8**, (1973) 1341.
- [5] G. Veneziano, Nuovo Cim. **A 57**, (1968) 190.
- [6] I. Niculescu et al., Phys. Rev. Letters **85**, (2000) 1182, 1186.
- [7] E.D. Bloom, E.J. Gilman, Phys. Rev. Letters **25**, (1970) 1149; Phys. Rev. **D 4**, (1971) 2901.
- [8] P. Freund, Phys. Rev. Letters **20**, (1968) 235; H. Harari, Phys. Rev. Letters **20**, (1968) 1395.
- [9] V. Rittenberg and H.R. Rubinstein, Nucl. Phys. **B 28**, (1971) 184;
- [10] G. Schierholz and M.G. Schmidt, Phys. Rev. **D 10**, (1974) 175.
- [11] A. De Rujula, H. Georgi, and H.D. Politzer, Ann. Phys. (N.Y.) **103**, (1977) 315; P. Stoler, Phys. Rev. **D 44**, (1991) 73; I. Afanasiev, C.E. Carlson, Ch. Wahlqvist, Phys. Rev. **D 62**, (2000) 074011; F.E. Close and N. Isgur, Phys. Letter **B 509**, (2001) 81; N. Isgur, S. Jeschonnek, W. Melnitchouk, and J.W. Van Orden, Phys. Rev. **D 64**, (2001) 054004.

- [12] P. Stoler, Phys. Rev. Letter **66**, (1991) 1003.
- [13] C.E. Carlson, N. Mukhopadhyay, Phys. Rev. **41**, (1990) 2343.
- [14] A.A. Logunov, L.D. Soloviov, A.N. Tavkhelidze Phys. Lett. **B 24**, (1967) 181; R. Dolen, D. Horn and C. Schmid Phys. Rev. **166**, (1968) 1768.
- [15] Marc Damashek and Frederick J. Gilman, Phys. Rev. **D 1**, (1970) 1319.
- [16] O. Nachtmann, Nucl. Phys. **B 63**, (1973) 237; *ibid* **B 78**, (1974) 455.
- [17] A. Bramon, E. Etim and M. Greco, Phys. Letters **B 41**, (1972) 609.
- [18] E. Etim, A. Malecki, Nuovo Cim **A 104**, (1991) 531.
- [19] A. Bugrij et al., Fortschritte der Physik, **21**, (1973) 427.
- [20] A. Degasperis and E. Predazzi, Nuovo Cim. **A 65**, (1970) 764.
- [21] R. Fiore, L. Jenkovszky, V. Magas, F. Paccanoni and A. Papa, hep-ph/0011035, Eur. Phys. J. **A 10**, (2001) 217.
- [22] M. Bertini, M. Giffon and E. Predazzi, Phys. Letters **B 349**, (1995) 561.
- [23] A. Capella, A. Kaidalov, C. Merino, J. Tran Thanh Van, Phys. Letters **B 337**, (1994) 358; L.P.A. Haakman, A. Kaidalov, J.H. Koch, Phys. Letters **365**, (1996) 411.
- [24] L. Jenkovszky, E. Martynov, F. Paccanoni, Padova preprint PFDPD 95/TH/21; P. Desgrolard, A.I. Lengyel, E.S. Martynov, EPJ **C 7**, (1999) 655.
- [25] M. Baillargeon and P. Nogueira, ‘Feynman diagram drawing made easy’, Nucl. Instr. and Meth. **A 389** (1997) 305.




Figure 9: Examples of least-squares fits to inclusive (e, e') data. The curves below the data are the individual resonance and nonresonant contributions. From [12].